NATtrol™ Epstein-Barr Virus Strain: B95-8

Catalog Number: NATEBV-0001 NATEBV-0002 NATEBV-0003 NATEBV-0004 NATEBV-0005 NATEBV-0006

PRODUCT DESCRIPTION:

NATtrol™ Epstein-Barr Virus (NATEBV-0001, NATEBV-0002, NATEBV-0003, NATEBV-0004, NATEBV-0005, and NATEBV-0006)* is formulated with purified, intact viral particles that have been chemically modified to render them non-infectious and refrigerator stable. Each vial contains 1.0 mL of NATtrol™ EBV in a proprietary matrix at the target concentration listed in Table 1.

*Pat.:http://www.zeptometrix.com/patent-information/

INTENDED USE:

NATtrol™ Epstein-Barr Virus is designed to evaluate the performance of nucleic acid tests for determination of the presence of EBV nucleic acid. NATtrol™ EBV enables laboratories to monitor test variation, lot-to-lot test kit performance, operator variation, and can provide assistance in identifying random or systemic error.

WARNINGS AND PRECAUTIONS:

- NATtrol™ inactivation was carried out on the EBV stock used to formulate this product. The inactivation was verified by the absence of viral growth of a surrogate virus in validated tissue culture based infectivity assays.
- This product contains inactivated microorganisms and materials of human and animal origin. Safe practices suggest that the controls be considered potentially infectious and to use Universal Precautions when
- Refer to CDC guidelines and local regulations for handling and disposal.
- The matrix used in the manufacture of this product is treated with 0.09% sodium azide. It was manufactured from Human Serum Albumin that has been tested and found to be non-reactive at the donor level for HIV-1/HIV-2 Antibody, HBsAg and HCV Antibody by FDA licensed donor screening test methods. All materials are also tested for HIV-1 and HCV by FDA approved Nucleic Acid Test (NAT) methods.
- Heat inactivated Fetal Bovine Serum used in the manufacture of this product meets applicable USDA requirements for abattoir sourced animals, traceability and country of origin. The materials were collected at USDA licensed establishments or legally imported from countries recognized by the USDA as negligible or controlled for risk for Bovine Spongiform Encephalopathy (BSE) and other exotic disease agents. Donor animals were inspected ante and post mortem at the abattoir as required by the USDA.
- Do not use past the expiration date on the label.
- To avoid cross-contamination, use separate pipette tips for all materials.

RECOMMENDED STORAGE:

NATtrol™ Epstein-Barr Virus should be stored at 2-8°C.

INSTRUCTIONS FOR USE:

- Mix vial vigorously for at least 5 secs.
- Process according to manufacturer's instructions for sample to result assays.
- Extract nucleic acid prior to use in downstream assays that are not sample to result.

LIMITATION:

- FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES
- Quality control materials should be used in accordance with local, state, federal, and accreditation requirements.
- This product is not intended to replace the manufacturer's controls provided with the assay.

EXPECTED RESULTS:

- Each laboratory must evaluate the product and establish their own acceptance criteria.
- The table shown below is for informational purposes only.

TABLE 1:

Catalog Number	Target Concentration (copies/mL) ^{1,2}	
NATEBV-0001	100	
NATEBV-0002	500	
NATEBV-0003	1,000	
NATEBV-0004	10,000	
NATEBV-0005	50,000	
NATEBV-0006	100,000	

¹Quantitation based on in-house real time PCR assay targeting the *LMP-2B* gene region.

ZMC-PI-Revision Effective

I-0031		Datcii Code	
n: 08	RUO	For Research Use Only	1
e Date: 02/09/2022	-	Manufacturer	

²Conversion factor: 1 copy = 1.62 IU. Based on internal testing of the 1st WHO International Standard for Epstein-Barr Virus for Nucleic acid Amplification Techniques (NIBSC code: 09/260)